



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

October 26, 2018

Dr. Chris Hofelt DABT
Product Registration Manager
BASF Corporation, Agricultural Products
26 Davis Drive, P.O. Box 13528
Research Triangle Park, NC 27709-3528

Subject: Label Amendment – Updates to Resistance-Management Labeling
Statements per PR Notices 2001-5 & 2017-1
Product Name: Nealta® Miticide
EPA Registration Number: 7969-336
Application Date: October 19, 2018
Decision Number: 545228

Dear Dr. Hofelt:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Page 2 of 2
EPA Reg. No. 7969-336
Decision No. 545228

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Mr. Carmen J. Rodia, Jr. by phone at (703) 306-0327, or via email at Rodia.Carmen@epa.gov.

Sincerely,

A handwritten signature in dark ink, appearing to read 'Gebken', written in a cursive style.

Richard Gebken
Product Manager 10
Invertebrate & Vertebrate Branch 2
Registration Division (7505P)
Office of Pesticide Programs

Enclosure: Stamped "Accepted" Master Label, dated October 26, 2018



We create chemistry

Cyflumetofen

Group

25

Acaricide

ACCEPTED

10/26/2018

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under

EPA Reg. No. 7969-336

Nealta®

Miticide

**For control of mites in citrus, grapes, pome fruits, strawberries,
tomatoes, and tree nuts**

Active Ingredient:

cyflumetofen: (2-methoxyethyl(*RS*)-2-(4-*tert*-butylphenyl)-2-cyano-3-oxo-3-
(α,α,α -trifluoro-*o*-tolyl)propionate) 18.7%

Other Ingredients: 81.3%

Total: 100.0%

Nealta® miticide contains 1.67 pounds of cyflumetofen per gallon.

EPA Reg. No. 7969-336

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN

CAUTION/PRECAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en
detalle. (If you do not understand this label, find someone to explain it to you in detail.)

See inside for complete **First Aid, Precautionary Statements, Directions For Use,**
Conditions of Sale and Warranty, and state-specific crop and/or use site restrictions.

In case of an emergency endangering life or property involving this product,
call day or night 1-800-832-HELP (4357).

Net Contents:

BASF Corporation
26 Davis Drive, Research Triangle Park, NC 27709

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to by a poison control center or doctor. • DO NOT give anything to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth to mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after first 5 minutes; then continue rinsing eyes. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), you may contact BASF Corporation at 1-800-832-HELP (4357), twenty-four (24) hours per day seven (7) days per week.</p>	

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION. Harmful if swallowed, absorbed through skin, or inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing spray mist.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber (includes natural rubber blends and laminates) ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, or viton ≥ 14 mils
- Shoes plus socks

Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

DO NOT apply directly to water, areas where surface water is present, or intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS ON ALL CROPS

- Apply this product only as specified the EPA approved label and any supplemental labeling.
- **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- **DO NOT** exceed the maximum seasonal use rate, the maximum rate per application, or the total number of **Nealta® miticide** applications per season as stated in the **Crop-specific Application Instructions** table.
- Preharvest Interval (PHI) restrictions are also included in this table.
- **DO NOT** apply through any type of irrigation system.
- **DO NOT** apply if rain is expected before the spray is dry for 1 hour. **Nealta** is rainfast 1 hour after drying.
- **DO NOT** apply **Nealta** via aerial application.
- **NOT** for use in greenhouses.
- For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber (includes natural rubber blends and laminates) ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, or viton ≥ 14 mils
- Shoes plus socks

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Keep this product in its tightly closed original container. Store in a cool, dry (preferably locked) area that is inaccessible to children and animals. **DO NOT** store near food or feed.

Pesticide Disposal

Wastes resulting from using this product may be disposed of on-site or at an approved waste disposal facility. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representatives at the nearest EPA Regional Office for guidance.

Container Handling

Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Triple rinse containers small enough to shake (capacity ≤ 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

In Case of Emergency

In case of large-scale spill of this product, call:

- CHEMTREC 1-800-424-9300
- BASF Corporation 1-800-832-HELP (4357)

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment
- Your local poison control center (hospital)
- BASF Corporation 1-800-832-HELP (4357)

Steps to take if this material is released or spilled:

- In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to label.
- Dike and contain the spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal.
- Remove contaminated clothing and wash affected skin areas with soap and water.
- Wash clothing before reuse.
- Keep the spill out of all sewers and open bodies of water.

Product Information

Nealta® miticide is a suspension concentrate formulation containing the active ingredient cyflumetofen, a member of the beta-ketonitrile class of chemistry. When used as directed, **Nealta** provides knockdown and residual control of tetranychid mites on the crops listed on this label.

Nealta is a highly active contact miticide on egg, nymph, and adult stage tetranychid mites. Because **Nealta** is not systemic and has no translaminar activity, thorough coverage of plant surfaces is necessary for effective control.

Nealta is **not** effective on non-tetranychid mites such as broad mite, flat mite, or rust mite. See **Pests Controlled** list for mites controlled.

Nealta is compatible with most biological control products.

Because **Nealta** is compatible with beneficial arthropods, it can be used effectively in Integrated Pest Management (IPM) and resistance management programs. **Nealta** does not have a negative impact on field populations of the following beneficial insects:

Beneficial Insects

Common Name	Scientific Name
Common lacewing	<i>Chrysopa carnea</i>
Insidious flower bug	<i>Orius insidiosus</i>
Predatory mite	<i>Amblyseius fallacis</i>
	<i>Phytoseiulus persimilis</i>
	<i>Typhlodromus pyri</i>
	<i>Zetzellia mali</i>
Seven-spotted lady beetle	<i>Coccinella septempunctata</i>
Six-spotted thrips	<i>Scolothrips sexmaculatus</i>
Spider mite destroyer	<i>Stethorus punctillum</i>
Western predatory mite	<i>Galendromus occidentalis</i>

Nealta is not systemic or translaminar; therefore, **thorough coverage of plant surfaces is important for optimum performance.**

Resistance-Management Recommendations

For resistance management, please note that cyflumetofen, the active ingredient in **Nealta**, belongs to the group of respiration inhibitors classified by the Insecticide Resistance Action Committee (IRAC) as target-site-of-action **Group 25** acaricide, a **Mitochondria complex II electron transport inhibitor**. Any mite population may contain individuals naturally resistant to **Nealta** and other **Group 25** acaricides. The resistant individuals may dominate the mite population if this group of acaricides are used repeatedly in the same fields. Repeated use of miticides with similar modes of action can lead to the buildup of resistant strains of mites.

- **DO NOT** exceed the maximum seasonal use rate or the total number of applications of **Nealta** per season.
- To reduce the potential for developing mite resistance, rotate to a miticide with a different mode of action.
- Monitor treated mite populations for resistance development.
- Read product label before applying any miticide and follow label directions.
- Use tank mixtures with acaricides from a different group that are equally effective on the target pest when such use is permitted. **DO NOT** rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity.
 - Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Contact your local extension specialist, certified crop advisor, and/or BASF representative for additional resistance management or IPM recommendations.

Plant Tolerance

Plant tolerance to **Nealta® miticide** has been found to be acceptable in research trials for the crops listed on this label. However, due to the large number of plant species and their associated varieties or cultivars, and due to variable growing conditions, it is impossible to test every plant for tolerance to this product.

Neither the manufacturer or seller has determined whether this product can be used without injury on all varieties of plants. It is the responsibility of the user to determine if this product can be used without causing plant injury prior to commercial use. In a small test area, test the specified rates on plants for phytotoxicity prior to large-scale use. The end user assumes all risks arising from application of **Nealta** in a manner inconsistent with the label requirements.

Pests Controlled

When used according to label instructions, **Nealta** is effective in controlling the following tetranychid mites:

Common Name	Scientific Name
Banks grass mite	<i>Oligonychus pratensis</i>
Brown almond mite	<i>Bryobia rubrioculus</i>
Brown wheat mite	<i>Petrobia lateens</i>
Carmine mite	<i>Tetranychus cinnabarinus</i>
Citrus red mite	<i>Panonychus citri</i>
European red mite	<i>Panonychus ulmi</i>
Spider mite, McDaniel	<i>Tetranychus mcdanieli</i>
Spider mite, Pacific	<i>Tetranychus pacificus</i>
Spider mite, spruce	<i>Oligonychus ununguis</i>
Spider mite, strawberry	<i>Tetranychus turkestanii</i>
Spider mite, two-spotted	<i>Tetranychus urticae</i>
Spider mite, Willamette	<i>Eotetranychus willamettei</i>
Spider mite, Yuma	<i>Eotetranychus yumensis</i>
Texas citrus mite	<i>Eutetranychus banksi</i>

Application Instructions

For maximum effectiveness, apply **Nealta** at the first sign of mites before the population increases. Application should be timed to coincide with locally recommended treatment threshold levels in developing mite populations.

Apply to foliage using properly calibrated ground sprayers (refer to **Crop-specific Application Instructions** table). Thorough and uniform spray coverage of foliage, with direct contact to target pest, is required for effective control. Applications made at lower volumes may result in less than thorough coverage which can lead to slower activity and/or less control than those made with higher spray volumes. Consult with your local or state extension personnel for advice on miticide use and selection.

Nealta is rainfast one (1) hour after an application has dried.

Apply Nealta in sufficient volume of water to ensure thorough coverage of foliage. For thorough coverage, use higher water volume.

Apply **Nealta** in minimum water volume per acre as indicated in **Crop-specific Application Instructions** table. Because **Nealta** is not systemic or translaminar, thorough coverage of plant surfaces is necessary for effective control. Applying **Nealta** at spray volumes lower than directed can make it harder to obtain thorough crop coverage and may reduce performance.

Adding an oil or adjuvant with spreading properties to the **Nealta** spray solution will provide improved plant coverage and optimum performance of **Nealta**. Test the safety and compatibility of all oils and adjuvants before use. Always read and follow the specific oil or adjuvant label using the proper concentration to avoid plant injury.

Controlling Spray Drift

Variable Winds. DO NOT apply in variable wind conditions.

Wind Direction. Wind direction (e.g. away from nontarget areas) must be based on the average direction (not instantaneous).

Wind Speed. Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size, canopy, and equipment specifications determine drift potential at any given wind speed. **DO NOT** apply when winds are greater than 15 mph.

Inversion Restriction. DO NOT make applications during temperature inversions. Drift potential is high during temperature inversions. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light-to-no wind.

Sensitive Areas. To ensure the protection of threatened or endangered species, it is important to maintain spray drift loadings below levels of concern for any area adjacent to the application site that is not excluded as possible habitat for these organisms. The following restrictions apply:

- **Buffer distance** - 15 feet
- **Release height** - 20 inches
- **Droplet size** - A combination of spray nozzles and appropriate pressure must be selected to provide ASABE standard S571.1 droplet size category of fine ($DV_{0.5}$ of ≥ 180 microns) or coarser than fine.
- **Maximum wind speed** - 15 mph

Cleaning Spray Equipment

Before application, start with clean, well-maintained application equipment. Following spray application, thoroughly clean all application equipment. Drain application equipment of any excess product. Thoroughly rinse application equipment and flush hoses, boom, and nozzles with clean water. Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. **DO NOT** clean equipment near wells, water sources, or desirable vegetation. Dispose of waste rinse-water in accordance with local regulations. Spraying equipment must be cleaned thoroughly before and after applying this product, particularly if a product with the potential to injure crops was used before **Nealta® miticide**.

Additives and Tank Mixing Information

Shake container well before use.

Nealta can be tank mixed with most recommended fungicides, insecticides, liquid fertilizers, biological control products, adjuvants, and additives.

Under some conditions, the use of additives or adjuvants can improve the performance of **Nealta**. However, all varieties and cultivars have not been tested with all possible tank mix combinations. Local conditions can also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, or crop injury can result from mixing **Nealta** with other products.

Before using any tank mix (fungicides, insecticides, liquid fertilizers, biological control products, adjuvants, and additives), test the combination on a small portion of the crop to be treated (including plant cultivars) to ensure that a phytotoxic response will not occur as a result of application. Evaluate for crop response 3 to 7 days before making an application to the entire crop.

When an adjuvant is to be used with this product, BASF recommends the use of a Chemical Producers and Distributors Association certified adjuvant.

Consult a BASF representative or local agricultural authorities for more information concerning additives.

Mixing Order

1. **Water** - Begin by agitating a thoroughly clean sprayer tank 3/4 full of clean water.
2. **Agitation** - Maintain constant agitation throughout mixing and application.
3. **Inductor** - If an inductor is used, rinse it thoroughly after each component has been added.
4. **Products in PVA bags** - Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5. **Water-dispersible products** (such as **Nealta**, dry flowables, wettable powders, other suspension concentrates, or suspo-emulsions)
6. **Water-soluble products**
7. **Emulsifiable concentrates** (such as oil concentrates when applicable)
8. **Water-soluble additives** (such as AMS or UAN when applicable)
9. **Remaining quantity of water**

Make sure each component is thoroughly mixed and suspended before adding tank mix partners. Maintain constant agitation during application.

Crop-specific Application Instructions

Crop	Pest	Product Use Rate per Application (fl ozs/A)	Minimum Time from Application to Harvest (PHI) days
Citrus Fruit Group Calamondin Citrus citron Citrus hybrids Chironja Grapefruit Kumquat Lemon Lime Mandarin Orange (sour and sweet) Pummelo Satsuma Tangelo Tangerine Tangor	See Pests Controlled under Product Information section	13.5 to 13.7	7

For knockdown and residual control of mites in citrus fruit, apply **Nealta® miticide** at first sign of infestation.

DO NOT make **Nealta** applications at intervals shorter than 14 days.

DO NOT make more than 2 applications per season per crop.

For maximum effectiveness, apply **Nealta** at the first sign of mites, and before the population increases. Application should be timed to coincide with locally recommended treatment threshold levels in developing mite populations.

Apply **Nealta** in at least 100 gallons of spray solution per acre; for **small nonbearing trees**, apply in at least 50 gallons of spray solution per acre. Applying **Nealta** at spray volumes lower than directed can make it harder to obtain thorough crop coverage and may reduce performance.

Resistance Management. DO NOT make more than 1 **Nealta** application before using an effective miticide with a different mode of action.

DO NOT apply more than 27.4 fl ozs/A (0.36 lbs cyflumetofen/A) per season per crop.

(continued)

Crop-specific Application Instructions *(continued)*

Crop	Pest	Product Use Rate per Application (fl ozs/A)	Minimum Time from Application to Harvest (PHI) days
Grapes	See Pests Controlled under Product Information section	13.5 to 13.7	14

For knockdown and residual control of mites in grapes, apply **Nealta® miticide** at first sign of infestation.

DO NOT make **Nealta** applications at intervals shorter than 14 days.

DO NOT make more than 2 applications per season per crop.

For maximum effectiveness, apply **Nealta** at the first sign of mites, and before the population increases. Application should be timed to coincide with locally recommended treatment threshold levels in developing mite populations.

Apply **Nealta** in at least 50 gallons of spray solution per acre. Applying **Nealta** at spray volumes lower than directed can make it harder to obtain thorough crop coverage and may reduce performance.

Resistance Management. DO NOT make more than 1 **Nealta** application before using an effective miticide with a different mode of action.

DO NOT apply more than 27.4 fl ozs/A (0.36 lbs cyflumetofen/A) per season per crop.

Crop	Pest	Product Use Rate per Application (fl ozs/A)	Minimum Time from Application to Harvest (PHI) days
Pome Fruit Group Apple Crabapple Loquat Mayhaw Oriental pear Pear Quince	See Pests Controlled under Product Information section	13.5 to 13.7	7

For knockdown and residual control of mites in pome fruit, apply **Nealta** at first sign of infestation.

DO NOT make **Nealta** applications at intervals shorter than 14 days.

DO NOT make more than 2 applications per season per crop.

For maximum effectiveness, apply **Nealta** at the first sign of mites, and before the population increases. Application should be timed to coincide with locally recommended treatment threshold levels in developing mite populations.

Apply **Nealta** in at least 100 gallons of spray solution per acre; for **small nonbearing trees**, apply in at least 50 gallons of spray solution per acre. Applying **Nealta** at spray volumes lower than directed can make it harder to obtain thorough crop coverage and may reduce performance.

Resistance Management. DO NOT make more than 1 **Nealta** application before using an effective miticide with a different mode of action.

DO NOT apply more than 27.4 fl ozs/A (0.36 lbs cyflumetofen/A) per season per crop.

(continued)

Crop-specific Application Instructions *(continued)*

Crop	Pest	Product Use Rate per Application (fl ozs/A)	Minimum Time from Application to Harvest (PHI) days
Strawberry	See Pests Controlled under Product Information section	13.5 to 13.7	1

For knockdown and residual control of mites in strawberries, apply **Nealta® miticide** at first sign of infestation.

DO NOT make **Nealta** applications at intervals shorter than 14 days.

DO NOT make more than 2 applications per season per crop.

For maximum effectiveness, apply **Nealta** at the first sign of mites, and before the population increases. Application should be timed to coincide with locally recommended treatment threshold levels in developing mite populations.

Apply **Nealta** in at least 50 gallons of spray solution per acre. Applying **Nealta** at spray volumes lower than directed can make it harder to obtain thorough crop coverage and may reduce performance.

Resistance Management. DO NOT make more than 1 **Nealta** application before using an effective miticide with a different mode of action.

DO NOT apply more than 27.4 fl ozs/A (0.36 lbs cyflumetofen/A) per season per crop.

Crop	Pest	Product Use Rate per Application (fl ozs/A)	Minimum Time from Application to Harvest (PHI) days
Tomato	See Pests Controlled under Product Information section	13.5 to 13.7	3

For knockdown and residual control of mites in tomatoes, apply **Nealta** at first sign of infestation.

DO NOT make **Nealta** applications at intervals shorter than 14 days.

DO NOT make more than 2 applications per season per crop.

For maximum effectiveness, apply **Nealta** at the first sign of mites, and before the population increases. Application should be timed to coincide with locally recommended treatment threshold levels in developing mite populations.

Apply **Nealta** in at least 50 gallons of spray solution per acre. Applying **Nealta** at spray volumes lower than directed can make it harder to obtain thorough crop coverage and may reduce performance.

Resistance Management. DO NOT make more than 1 **Nealta** application before using an effective miticide with a different mode of action.

DO NOT apply more than 27.4 fl ozs/A (0.36 lbs cyflumetofen/A) per season per crop.

(continued)

Crop-specific Application Instructions *(continued)*

Crop	Pest	Product Use Rate per Application (fl ozs/A)	Minimum Time from Application to Harvest (PHI) days
Tree Nuts Group Almonds Beech nut Brazil nut Butternut Cashew Chestnut Chinquapin Filbert Hickory nut Macadamia nut Pecan Pistachio Walnuts (black and English)	See Pests Controlled under Product Information section	13.5 to 13.7	7

For knockdown and residual control of mites in tree nuts, apply **Nealta® miticide** at first sign of infestation.

DO NOT make **Nealta** applications at intervals shorter than 14 days.

DO NOT make more than 2 applications per season per crop.

For maximum effectiveness, apply **Nealta** at the first sign of mites, and before the population increases. Application should be timed to coincide with locally recommended treatment threshold levels in developing mite populations.

Apply **Nealta** in at least 100 gallons of spray solution per acre; for **small nonbearing trees**, apply in at least 50 gallons of spray solution per acre. Applying **Nealta** at spray volumes lower than directed can make it harder to obtain thorough crop coverage and may reduce performance.

Resistance Management. DO NOT make more than 1 **Nealta** application before using an effective miticide with a different mode of action.

DO NOT apply more than 27.4 fl ozs/A (0.36 lbs cyflumetofen/A) per season per crop.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

Nealta is a registered trademark of BASF.

© 2018 BASF Corporation
All rights reserved.

007969-00336.20181002b.**NVA 2018-04-390-0081**
Supersedes: NVA 2016-04-390-0112

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709


We create chemistry